Covert Administration of Oral Medication Plan



This form is intended to collate all relevant signatures and requirements onto one place to allow ease of reviews, it is not intended to replace support documentation and an understanding of the legal requirements around safe Covert administration practices

In line with good practice, capacity should be reviewed as close to six monthly as possible, ideally as part of the GP enhanced service review, or earlier if there has been a change in their condition and/or their medication

In order to determine whether a patient has mental capacity, a test of mental capacity must be carried out in accordance with that set out in the Mental Capacity Act 2005 and MCA code of practice 2007 and should only consider the residents capacity around their understanding of the medication they have been prescribed (see below)

A patient will be considered to lack mental capacity if he or she is unable to:							Υ	N	
Understand in simple langua	age what the treatment i	s, its purpose and why it is bei	ing proposed.						
Understand its principle ben	efits, risks and alternativ	/es.							
Understand in broad terms what will be the consequences of not receiving the proposed treatment.									
Retain the information long	enough to make an effe	ctive decision, which will be fre	ee from pressure.						
Communicate that decision.									
If the answer to any of the qu	estions is No then the re	esident does not have mental o	capacity.						
Residents Name			Date	of Birth		Date of assess	ment		
Reason for presuming ment	al incapacity								
Has the reason been record	ed in the proposed man	agement plan and a Best Inter	rest form completed					Υ	N
Assessors name				Signature					
The discussion to covertly ac	Iminister medication has	s been discussed and agreed v	with;						
	Signature	Review 1 Date	Review 2 Date	Review 3 Date			Review 4 Date		
GP									
Care Home Nurse									
Next of kin or Advocate									
Pharmacist consulted									

Before crushing etc. any medication please consider liquid or dispersible formulation alternatives

Medication to be administered covertly	Method of administration i.e. crushed/sprinkle				
Always put into the smallest amount of food or liquid possible to unsure all the medication is taken	·				
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					
11					
12					
13					
14					
15					
16					
17					
18					

If any of the medications listed could affect mental capacity, there should be a referral to the DOLS's team