

## Patient Information Undergoing Percutaneous Nephrostomy

**Directorate of Clinical Radiology**

### **Introduction**

This leaflet tells you about the procedure known as percutaneous nephrostomy, explains what is involved and what the possible risks are. It is not meant to replace informed discussion between you and your doctor but can act as a starting point for such a discussion.

If the percutaneous nephrostomy is being done as a pre-planned procedure, then you should have plenty of time to discuss the situation with your consultant and the radiologist who will be doing the percutaneous nephrostomy, and perhaps even your own GP. If you need the nephrostomy as an emergency, then there may be less time for discussion, **but none the less you should have had sufficient explanation before you sign the consent form.**

### **What is a percutaneous nephrostomy?**

The urine from a normal kidney drains through a narrow, muscular tube, the urethra, into the bladder. When that tube becomes blocked, for example by a stone or a blood clot, the kidney can rapidly become affected, especially if there is infection present as well.

While an operation may become necessary, it is also possible to relieve the blockage by inserting a fine plastic tube, called a catheter, through the skin, into the kidney, under local anaesthetic. This catheter then allows the urine to drain from the kidney into a collecting bag, outside the body. This procedure is called a percutaneous (meaning through the skin) nephrostomy (a tube put into the kidney).

### **Why do I need a percutaneous nephrostomy?**

Other tests will have shown that the tube leading from your kidney to the bladder has become blocked. However, it may not be obvious what the cause of the blockage is. If left untreated, your kidney may become damaged.

## **Who has made the decision?**

The doctors in charge of your case, and the radiologist doing the percutaneous nephrostomy will have discussed the situation, and feel that this is the best treatment option. However, you will also have the opportunity for your opinion to be considered, and if, after discussion with your doctors, you do not want the procedure carried out, then you can decide against it.

## **Who will be doing the percutaneous nephrostomy?**

A specially trained doctor called a radiologist. Radiologists have special expertise in using x-ray and scanning equipment, and also in interpreting the images produced. They need to look at these images while carrying out the procedure. Consequently, radiologists are the best-trained people to insert needles and fine tubes into the body, through the skin, and place them correctly.

## **Medication**

The Radiologist will need to know all medication that you are taking, including anti-platelet drugs and anticoagulants. These are medicines that affect the way your blood clots.

Listed below are some drugs that the doctor will need to be informed about if you have been prescribed them:

Warfarin	Aspirin	Enoxaparin	Dalteparin
Clopidogrel	Tinzaparin	Bemiparin	Fondaparinux
Danaparoid	Dipyridamole	Coumarins	Acenocoumarol
Phenindione	Lepirudin	GP IIb/IIIa inhibitors	
Bivalirudin	Argatroban	Dabigatran	Rivaroxaban
Abciximab	Eptifibatide	Tirofiban	Plasugrel

The Radiologist will need to know if you have a hereditary bleeding abnormality or abnormal bleeding history after minor procedures such as dental extraction and further tests to assess your blood may be needed.

## **Where will the procedure take place?**

Generally, in the x-ray department, in a special "screening" room, which is adapted for specialised procedures. It may be done in a C.T. scanner or an operating theatre, using mobile x-ray equipment or a portable ultrasound scanner.

## **How do I prepare for percutaneous nephrostomy?**

You need to be an in-patient in the hospital.

You will probably be asked not to eat for four hours beforehand, though you may be allowed to drink some water.

You may receive a sedative to relieve anxiety, as well as an antibiotic.

You will be asked to put on a hospital gown.

If you have any allergies, you **must** let your doctor know. If you have previously reacted to intravenous contrast medium, the dye used for kidney x-rays and CT scanning, then you must also tell your doctor about this.

## **What actually happens during a percutaneous nephrostomy?**

You will lie on the x-ray table, generally flat on your stomach, or nearly flat. You need to have a needle put into a vein in your arm, so that the radiologist can give you a sedative or painkillers. Once in place, this needle does not cause any pain. You will also have a monitoring device attached to your chest and finger, and will probably receive oxygen through small tubes in your nose. The radiologist will keep everything as sterile as possible, and may wear a theatre gown and operating gloves. Your skin will be cleaned with antiseptic, and then most of the rest of your body covered with a theatre towel.

The radiologist will use the x-ray equipment or the ultrasound machine to decide on the most suitable point for inserting the fine plastic tube (catheter), usually in your back, just below your bottom rib. Then your skin will be anaesthetised with local anaesthetic, and a fine needle inserted into the kidney. When the radiologist is sure that the needle is in a satisfactory position, a guide wire will be placed into the kidney, through the needle, which then enables the plastic catheter to be positioned correctly. This catheter will then be fixed to the skin surface, and attached to a drainage bag.

## **Will it hurt?**

Unfortunately, it may hurt a little, for a very short period of time, but any pain you have should be controlled with painkillers.

When the local anaesthetic is injected, it will sting to start with, but this soon wears off, and the skin and deeper tissues should then feel numb. Later, you may be aware of the needle and then the catheter passing into the kidney, and sometimes this is painful, especially if the kidney was sore to start with.

There will be a nurse, or another member of clinical staff, standing next to you and looking after you.

If the procedure does become painful for you, then they will be able to arrange for you to have more painkillers through the needle in your arm. Generally, placing the catheter in the kidney only takes a short time, and once in place it should not hurt at all.

## **How long will the catheter stay in, and what happens next?**

These are questions, which only the doctors looking after you can answer. It may only need to stay in a short time, for example while a stone passes naturally, or it may need to stay in for a much longer period, to allow a more permanent solution for the blockage to be organised. Taking the catheter out does not hurt at all.

## **How long will it take?**

Every patient's situation is different, and it is not always easy to predict how complex or how straightforward the procedure will be. It may be over in 20 minutes, or very occasionally it may take longer than 90 minutes. As a guide, expect to be in the x-ray department for about an hour altogether.

## **What happens afterwards?**

You will be taken back to your ward on a trolley. Nurses on the ward will carry out routine observations, such as taking your pulse and blood pressure, to make sure that there are no problems. You will generally stay in bed for a few hours, until you have recovered.

The drainage catheter stays in place in your body for the time being, and will be attached to a collection bag. You will be able to carry on a normal life with the catheter in place.

However, it is important that you try not to make any sudden movements, for example getting up out of a chair, without remembering about the bag, and making sure that it can move freely with you. The bag needs to be emptied fairly frequently, so that it does not become too heavy, but the nurses will want to measure the amount in it each time.

### **Are there any risks or complications?**

Percutaneous nephrostomy is a very safe procedure, but there are some risks and complications that can arise, as with any medical treatment. Perhaps the biggest problem is being unable to place the drainage tube satisfactorily in the kidney. If this happens, a surgeon will arrange another method of overcoming the blockage, which may involve surgery.

Sometimes there is a leak of urine from the kidney, resulting in a small collection of fluid inside the abdomen. If this becomes a large collection, it may require draining.

There may be slight bleeding from the kidney. On very rare occasions, this may become severe, and require a surgical operation or another radiological procedure to stop it. Occasionally there may be infection in the kidney, or in the space around it. This can generally be treated satisfactorily with antibiotics. Risk of death under 1:1000

Despite these possible complications, the procedure is normally very safe, and will almost certainly result in a great improvement in your medical condition. Very occasionally, an operation is required, but if the percutaneous nephrostomy had not been attempted, then this operation would have been necessary anyway.

### **Further Information**

Some of your questions should have been answered by this leaflet, but remember that this is only a starting point for discussion about your treatment with the doctors looking after you. **Make sure you are satisfied that you have received enough information about the procedure, before you sign the consent form.**

Percutaneous nephrostomy is considered a very safe procedure, designed to save you having a larger operation. There are some slight risks and possible complications involved, and although it is difficult to say exactly how often these occur, they are generally minor and do not happen very often.

**If you have any questions or concerns, please find below contact numbers for the Urology wards and Urology nurses:-**

**Urology Ward D2 East**

Tel. No:- 01633 - 234043 / 234044  
24 hours

**Urology Outpatients Department**

Tel. No:- 01633 - 234979  
Monday – Friday office hours

**Urology Day Ward**

Tel. No:- 01633 – 656378 / 656377  
Monday – Friday office hours

**Uro-Oncology Clinical Nurse Specialists**

Julie Simpson

Sian Lewis

Louise Broadway

Kathleen Connor-Middleton

Tel. No: 01633 – 238976 (will return voicemails when not on clinical duties)

Monday – Friday office hours

**Urology Nurse Practitioner**

Laura Reynolds

Leanne Denning

Tel. No:- 01633 – 234758

Monday – Friday office hours

**Surgery/Pre-admission queries**

Tel. No:- 01633 – 238568

Monday – Friday office hours

Modified with thanks from ©The Royal College of Radiologists, July 2000

This leaflet has been prepared by the British Society of Interventional Radiology (BSIR) and the Clinical Radiology Patients' Liaison Group (CRPLG) of the Royal College of Radiologists.

**“This document is available in Welsh /  
Mae'r ddogfen hon ar gael yn Gymraeg”.**