



GIG  
CYMRU  
NHS  
WALES

Bwrdd Iechyd Prifysgol  
Aneurin Bevan  
University Health Board

## Discharge advice following attendance with a paediatric (children's) clavicle fracture

### Trauma and Orthopaedics

- Your child has fractured their Clavicle (Collar Bone)
- This type of fracture is common in children
- This type of fracture heals well – the only treatments required are painkillers and a sling
- We would expect the collar bone to be painful for 4 – 6 weeks
- Your child may find it more comfortable to sleep sitting upright for a few days after the injury
- The shoulder and arm can be moved out of the sling as comfort allows. This will usually be about 2 weeks after the injury but can be sooner if comfortable
- The “bump” over the fracture is quite normal and is produced by healing bone. It may take up to one year to disappear. If your child is older than ten years a small bump may remain
- Your child may return to sports such as swimming as soon as comfortable, but should avoid contact sports such as football. Rugby and basketball for six weeks



- If you are still experiencing significant symptoms after several months, please phone the fracture clinic helpline as listed below for further advice

Should you have any worries or concerns following discharge from the hospital, please contact: 01633 234403/234404. This is a Monday – Friday service and we will try to get back to you as soon as possible. However, if it is an emergency please go back to A&E for further advice/treatment.

**This document is available in Welsh /  
Mae'r ddogfen hon ar gael yn Gymraeg**